

ON A THEOREM OF MACAULAY ON COLONS OF IDEALS*

J. K. VERMA

Dedicated to my teacher William Heinzer

ABSTRACT. Let R be a polynomial ring over a field of dimension n . Let $\mathfrak{m}$ be the maximal homogeneous ideal of R . Let I be a complete intersection homogeneous ideal of R minimally generated by polynomials of degrees $d_1, d_2, \dots, d_n$. F. S. Macaulay showed that for all integers $i = 0, 1, \dots, \delta + 1$, $I : \mathfrak{m}^i = I + \mathfrak{m}^{\delta+1-i}$ where $\delta = d_1 + d_2 + \dots + d_n - n$. We provide a modern proof of a generalization of this result to standard graded Gorenstein rings.

In §86 of his monograph [M] of 1916, F. S. Macaulay proved the following

Theorem 1.1. *Let R be the polynomial ring $k[X_1, X_2, \dots, X_n]$ over a field k . Let I be a height n ideal of R generated by homogeneous polynomials $f_1, f_2, \dots, f_n$ of degrees $d_1, d_2, \dots, d_n$ respectively. Set $\mathfrak{m} = (X_1, X_2, \dots, X_n)R$ and $\delta = d_1 + d_2 + \dots + d_n - n$. Then for all integers $i = 0, 1, \dots, \delta + 1$,*

$$I : \mathfrak{m}^i = I + \mathfrak{m}^{\delta+1-i}.$$

In his expository paper [G] P. Griffiths used Macaulay's theorem in Hodge theory of smooth hypersurfaces in projective space. He also provided a proof of this theorem by using the local duality theorem in the complex case involving the Grothendieck residue symbol. The objective of this note is to provide a modern algebraic proof which seems to be lacking in the literature. We will generalize Macaulay's theorem to $\mathfrak{m}$ -primary ideals in standard graded Gorenstein rings.

By taking the quotient of R by I , it is enough to prove the theorem for standard graded Artinian Gorenstein rings.

*Presented at the Third National Conference on Commutative Algebra and Algebraic Geometry, Indian Institute of Science, Bangalore, India, October 16-21, 2000.

A crucial result used in our proof is a theorem of Macaulay about Hilbert series of Gorenstein graded algebras. For a modern proof, see Corollary 4.4.6 of [BH]. This has been generalized to Gorenstein graded rings over Artinian rings in [JV]. We shall use the zero dimensional case of this. We recall the statement for the convenience of the reader. Let $\ell(\cdot)$ denote length.

Theorem 1.2. *Suppose that $S = \bigoplus_{n=0}^{\infty} S_n$ is a standard graded Artinian Gorenstein algebra over an Artinian local ring S_0 . Let $\delta = \max\{n \mid S_n \neq 0\}$. Then $\ell(S_i) = \ell(S_{\delta-i})$ for all $i = 0, 1, \dots, \delta$.*

We recall certain standard facts from [BH] about local cohomology. Let $R = \bigoplus_{n=0}^{\infty} R_n$ be a d -dimensional finitely generated standard graded algebra over a Noetherian local ring R_0 . Let $\mathfrak{m}$ denote the irrelevant ideal $\bigoplus_{n=1}^{\infty} R_n$. The a -invariant of R is defined by

$$a(R) = \max\{n \mid [H_{\mathfrak{m}}^d(R)]_n \neq 0\}.$$

If R is Cohen-Macaulay and f is a homogeneous nonzerodivisor of degree d then $a(R/fR) = a(R) + d$. If R is a polynomial ring over a field in n variables then $a(R) = -n$. Thus If $\mathbf{f} = (f_1, f_2, \dots, f_n)$ is a regular sequence in a polynomial ring with $\deg(f_i) = d_i$ for all $i = 1, 2, \dots, n$ then $a(R/\mathbf{f}) = d_1 + d_2 + \dots + d_n - n$. This explains the value of δ in Macaulay's theorem.

Theorem 1.3. *Let $(R, \mathfrak{m})$ be an Artinian Gorenstein local ring. Let I be an $\mathfrak{m}$ -primary ideal. Put $H(I, n) = \ell(I^n/I^{n+1})$. Let $\delta = \max\{i \mid I^i \neq 0\}$. Then $0 : I^i = I^{\delta+1-i}$ for all $i = 0, 1, \dots, \delta$ if and only if $H(I, i) = H(I, \delta - i)$ for all $i = 0, 1, \dots, \delta$.*

Proof. Since R is Gorenstein, it is the injective hull of its residue field. Hence by Matlis duality $\ell(\text{Hom}(R/I^i, R)) = \ell(R/I^i)$ for all i . Since $I^{\delta+1-i} \subseteq (0 : I^i)$ and $\ell(\text{Hom}(R/I^i, R)) = \ell(0 : I^i)$, we conclude that $0 : I^i = I^{\delta+1-i}$ for all $i = 0, 1, \dots, \delta$ if and only if $\ell(R/I^i) = \ell(I^{\delta+1-i})$ for all $i = 0, 1, \dots, \delta$ if and only if $H(I, 0) + \dots + H(I, i) = H(I, \delta - i) + \dots + H(I, \delta)$ for all $i = 0, 1, \dots, \delta$. It is easy to see that the last statement is equivalent to the condition $H(I, i) = H(I, \delta - i)$ for all $i = 0, 1, \dots, \delta$. □

Corollary 1.4. *Let R be a standard graded Artinian Gorenstein ring over an Artinian local ring $(S, \mathfrak{n})$. Let $\mathfrak{m}$ be the graded ideal of R generated by elements of positive degree*

with $\mathfrak{m}^\delta \neq 0$ and $\mathfrak{m}^{\delta+1} = 0$. Then for all $i = 0, 1, \dots, \delta$,

$$0 : \mathfrak{m}^i = \mathfrak{m}^{\delta+1-i}.$$

Proof. Since R is standard, the associated graded ring of R with respect to $\mathfrak{m}$ is R itself. Since R is Gorenstein, its Hilbert function is symmetric by [JV].

□

Example 1.5. We present an example due to U. Storch which shows that Macaulay's theorem is not valid for Gorenstein local rings. Let k be a field of characteristic 2. Let X and Y be indeterminates. Put $R = k[X, Y]/I$ where $I = (X^2 + Y^2, X^2 + XY + Y^3)$. Then R is a Gorenstein local ring. The Hilbert series $H(G, \lambda) = \sum_{i=0}^{\infty} \ell(\mathfrak{m}^i/\mathfrak{m}^{i+1})\lambda^i$ of the associated graded G of R with respect to its maximal ideal $\mathfrak{m}$ is given by

$$H(G, \lambda) = 1 + 2\lambda + \lambda^2 + \lambda^3.$$

Thus the Hilbert function is not symmetric. Hence Macaulay's theorem does not hold for R by Theorem 1.3.

Acknowledgement : The author thanks U. Storch and B. Singh for useful conversations.

REFERENCES

- [BH] W. Bruns and J. Herzog, **Cohen-Macaulay rings**, revised edition, Cambridge Studies in Advanced Mathematics, 39. Cambridge University Press, Cambridge, 1998.
- [G] P. A. Griffiths, Complex analysis and algebraic geometry, Bull. Amer. Math. Soc. (N.S.) 1 (1979), no. 4, 595–626.
- [JV] B. Johnston and J.K. Verma, On the length formula of Hoskin and Deligne and associated graded rings of two-dimensional regular local rings, Math. Proc. Cambridge Philos. Soc. 111 (1992), 423–432.
- [M] F. S. Macaulay, **The algebraic theory of modular systems**, Cambridge University Press (1916), reissued with an introduction by P. C. Roberts (1994).

DEPARTMENT OF MATHEMATICS, IIT BOMBAY, POWAI, MUMBAI, INDIA - 400076

E-mail address: jkv@math.iitb.ac.in